

SUMMER REVIEW PACKET

For students entering IB Math Studies I

Name:

The attached assignment covers prerequisites for math studies. In other words, these are objectives that you should have mastered in geometry and algebra 2. We will be testing on these objectives the first or second class of the year. The assignment is an opportunity for you to review and ensure that you have mastered the prerequisites.

Radicals:

To simplify means that 1) no radicand has a perfect square factor and
2) there is no radical in the denominator (rationalize).

Recall – the **Product Property** $\sqrt{ab} = \sqrt{a} \cdot \sqrt{b}$ and the **Quotient Property** $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$

Examples: Simplify $\sqrt{24} = \sqrt{4} \cdot \sqrt{6}$ find a perfect square factor
 $= 2\sqrt{6}$ simplify

Simplify $\sqrt{\frac{7}{2}} = \frac{\sqrt{7}}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}}$ split apart, then multiply by both the numerator and the
denominator by $\sqrt{2}$
 $= \frac{\sqrt{14}}{\sqrt{4}} = \frac{\sqrt{14}}{2}$ multiply straight across and simplify

If the denominator contains 2 terms –
multiply the numerator and the denominator by the *conjugate* of the denominator
The *conjugate* of $3 + \sqrt{2}$ is $3 - \sqrt{2}$ (the sign changes between the terms).

Simplify each of the following.

1. $\sqrt{32}$

2. $\sqrt{(2x)^8}$

3. $\sqrt[3]{-64}$

4. $\sqrt{49m^2n^8}$

5. $\sqrt{\frac{11}{9}}$

6. $\sqrt{60} \cdot \sqrt{105}$

7. $(\sqrt{5} - \sqrt{6})(\sqrt{5} + \sqrt{2})$

Rationalize.

8. $\frac{1}{\sqrt{2}}$

9. $\frac{2}{\sqrt{3}}$

10. $\frac{3}{2 - \sqrt{5}}$

Complex Numbers:Form of complex number : $a + bi$ Where a is the "real" part and bi is the "imaginary" partAlways make these substitutions $\sqrt{-1} = i$ and $i^2 = -1$

- To simplify: pull out the $\sqrt{-1}$ before performing any operation

Example: $\sqrt{-5} = \sqrt{-1} \cdot \sqrt{5}$ Pull out $\sqrt{-1}$ **Example:** $(i\sqrt{5})^2 = i\sqrt{5} \cdot i\sqrt{5}$ List twice
 $= i\sqrt{5}$ Make substitution $= i^2 \sqrt{25}$ Simplify
 $= (-1)(5) = -5$ Substitute

- Treat i like any other variable when +, -, \times , or \div (but always simplify $i^2 = -1$)

Example: $2i(3 + i) = 2(3i) + 2i(i)$ Distribute
 $= 6i + 2i^2$ Simplify
 $= 6i + 2(-1)$ Make substitution
 $= -2 + 6i$ Simplify and rewrite in complex form

- Since $i = \sqrt{-1}$, no answer can have an 'i' in the denominator **RATIONALIZE!!**

Simplify.

11. $\sqrt{-49}$

12. $6\sqrt{-12}$

13. $-6(2 - 8i) + 3(5 + 7i)$




14. $(3-4i)^2$

15. $(6-4i)(6+4i)$



Geometry:

Pythagorean Theorem (right triangles): $a^2 + b^2 = c^2$



Find the value of x .




19. A square has perimeter 12 cm. Find the length of the diagonal.

In $30^\circ - 60^\circ - 90^\circ$ triangles, Sides are in proportion $1 : \sqrt{3} : 2$	In $45^\circ - 45^\circ - 90^\circ$ triangles, Sides are in proportion $1 : 1 : \sqrt{2}$
	


Solve for x and y .



22.



23.



Equations of Lines:

Slope intercept form: $y = mx + b$

Vertical line: $x = a$ (slope is undefined)

Point-slope form: $y - y_1 = m(x - x_1)$

Horizontal line: $y = b$ (slope is 0)

Standard Form: $Ax + By = C$

Slope: $m = \frac{y_2 - y_1}{x_2 - x_1}$

24. State the slope and y-intercept of the linear equation: $5x - 4y = 8$.

25. Find the x-intercept and y-intercept of the equation: $2x - y = 5$

26. Write the equation in standard form: $y = 7x - 5$

Write the equation of the line in slope-intercept form with the following conditions:

28. slope = -5 and passes through the point $(-3, -8)$


27. passes through the points $(4, 3)$ and $(7, -2)$

29. x-intercept = 3 and y-intercept = 2


Graphing:

Graph each function, inequality, and / or system.


30. $3x - 4y = 12$




31. $\begin{cases} 2x + y = 4 \\ x - y = 2 \end{cases}$




32. $y < -4x - 2$



33. $y + 2 = |x + 1|$



34. $y + 4 = (x - 1)^2$



Vertex: _____

x-intercept(s): _____

y-intercept(s): _____

Systems of Equations:

$$3x + y = 6$$
$$2x - 2y = 4$$

Substitution:

Solve 1 equation for 1 variable.
Rearrange.
Plug into 2nd equation.
Solve for the other variable.

Then plug answer back into an original equation to solve for the 2nd variable.

$$y = 6 - 3x \quad \text{solve 1st equation for } y$$
$$2x - 2(6 - 3x) = 4 \quad \text{plug into 2nd equation}$$
$$2x - 12 + 6x = 4 \quad \text{distribute}$$
$$8x = 16 \quad \text{simplify}$$
$$x = 2$$

Elimination:

Find opposite coefficients for 1 variable.
Multiply equation(s) by constant(s).
Add equations together (lose 1 variable).
Solve for variable.

$$6x + 2y = 12 \quad \text{multiply 1st equation by 2}$$
$$2x - 2y = 4 \quad \text{coefficients of } y \text{ are opposite}$$
$$\hline 8x = 16 \quad \text{add}$$
$$x = 2 \quad \text{simplify}$$

$$3(2) + y = 6$$

Plug $x = 2$ back into original

$$6 + y = 6$$
$$y = 0$$

Solve each system of equations. Use any method.

35.
$$\begin{cases} 2x + y = 4 \\ 3x + 2y = 1 \end{cases}$$

36.
$$\begin{cases} 2x + y = 4 \\ 3x - y = 14 \end{cases}$$

Exponents:

TWO RULES OF ONE

1. $a^1 = a$

Any number raised to the power of one equals itself.

2. $1^a = 1$

One to any power is one.

ZERO RULE

3. $a^0 = 1$

Any nonzero number raised to the power of zero is one.

PRODUCT RULE

4. $a^m \cdot a^n = a^{m+n}$

When multiplying two powers that have the same base, add the exponents.

QUOTIENT RULE

5. $\frac{a^m}{a^n} = a^{m-n}$

When dividing two powers with the same base, subtract the exponents.

POWER RULE

6. $(a^m)^n = a^{mn}$

When a power is raised to another power, multiply the exponents.

NEGATIVE EXPONENTS

7. $a^{-n} = \frac{1}{a^n}$ and $\frac{1}{a^{-n}} = a^n$

Any nonzero number raised to a negative power equals its reciprocal raised to the opposite positive power.

Express each of the following in simplest form. Answers should not have any negative exponents.

37. $5a^0$

38. $\frac{3c}{c^{-1}}$

39. $\frac{2ef^{-1}}{e^{-1}}$

40. $\frac{(n^3 p^{-1})^2}{(np)^{-2}}$

Simplify.

41. $3m^2 \cdot 2m$

42. $(a^3)^2$

43. $(-b^3 c^4)^5$

44. $4m(3a^2 m)$

Polynomials:

To add / subtract polynomials, combine like terms.

$$\begin{aligned} \text{EX: } & 8x - 3y + 6 - (6y + 4x - 9) && \text{Distribute the negative through the parentheses.} \\ & = 8x - 3y + 6 - 6y - 4x + 9 && \text{Combine terms with similar variables.} \\ & = 8x - 4x - 3y - 6y + 6 + 9 \\ & = 4x - 9y + 15 \end{aligned}$$

Simplify.

45. $3x^3 + 9 + 7x^2 - x^3$

46. $7m - 6 - (2m + 5)$

To multiplying two binomials, use FOIL.

$$\begin{aligned} \text{EX: } & (3x - 2)(x + 4) && \text{Multiply the first, outer, inner, then last terms.} \\ & = 3x^2 + 12x - 2x - 8 && \text{Combine like terms.} \\ & = 3x^2 + 10x - 8 \end{aligned}$$

To multiplying two binomials, use FOIL.

$$\begin{aligned} \text{EX: } & (3x - 2)(x + 4) && \text{Multiply the first, outer, inner, then last terms.} \\ & = 3x^2 + 12x - 2x - 8 && \text{Combine like terms.} \\ & = 3x^2 + 10x - 8 \end{aligned}$$

Multiply.

47. $(3a + 1)(a - 2)$

48. $(s + 3)(s - 3)$

49. $(c - 5)^2$

50. $(5x + 7y)(5x - 7y)$

Factoring.

Follow these steps in order to factor polynomials.

STEP 1: Look for a GCF in ALL of the terms.

- a.) If you have one (other than 1) factor it out front.
 b.) If you don't have one, move on to STEP 2.

STEP 2: How many terms does the polynomial have?**2 Terms**a.) Is it difference of two squares? $a^2 - b^2 = (a + b)(a - b)$

EX: $x^2 - 25 = (x + 5)(x - 5)$

b.) Is it sum or difference of two cubes?

$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$

$a^3 + b^3 = (a + b)(a^2 - ab + b^2)$

EX: $m^3 + 64 = (m + 4)(m^2 - 4m + 16)$

$p^3 - 125 = (p - 5)(p^2 + 5p + 25)$

3 Terms

$x^2 + bx + c = (x + \quad)(x + \quad)$

Ex: $x^2 + 7x + 12 = (x + 3)(x + 4)$

$x^2 - bx + c = (x - \quad)(x - \quad)$

$x^2 - 5x + 4 = (x - 1)(x - 4)$

$x^2 + bx - c = (x - \quad)(x + \quad)$

$x^2 + 6x - 16 = (x - 2)(x + 8)$

$x^2 - bx - c = (x - \quad)(x + \quad)$

$x^2 - 2x - 24 = (x - 6)(x + 4)$

4 Terms -- Factor by Grouping

- a.) Pair up first two terms and last two terms
 b.) Factor out GCF of each pair of numbers.
 c.) Factor out front the parentheses that the terms have in common.
 d.) Put leftover terms in parentheses.

$$\begin{aligned} \text{Ex: } x^3 + 3x^2 + 9x + 27 &= (x^3 + 3x^2) + (9x + 27) \\ &= x^2(x + 3) + 9(x + 3) \\ &= (x + 3)(x^2 + 9) \end{aligned}$$

Factor completely.

51. $z^2 + 4z - 12$

52. $6 - 5x - x^2$

53. $2k^2 + 2k - 60$

54. $-10b^4 - 15b^2$

55. $9c^2 + 30c + 25$

56. $9m^2 - 4$

To solve quadratic equations, try to factor first and set each factor equal to zero. Solve for your variable. If the quadratic does NOT factor, use quadratic formula.

EX: $x^2 - 4x = 21$ *Set equal to zero FIRST.*
 $x^2 - 4x - 21 = 0$ *Now factor.*
 $(x + 3)(x - 7) = 0$ *Set each factor equal to zero.*
 $x + 3 = 0$ $x - 7 = 0$ *Solve each for x.*
 $x = -3$ $x = 7$

Solve each equation.


57. $x^2 - 4x - 12 = 0$

58. $x^2 + 25 = 10x$


59. $x^2 - 14x + 40 = 0$

DISCRIMINANT: The number under the radical in the quadratic formula ($b^2 - 4ac$) can tell you what kinds of roots you will have.


IF $b^2 - 4ac > 0$ you will have TWO real roots.
(touches x-axis twice)



IF $b^2 - 4ac = 0$ you will have ONE real root
(touches the x-axis once)



IF $b^2 - 4ac < 0$ you will have TWO imaginary roots.
(Graph does not cross the x-axis)



QUADRATIC FORMULA – allows you to solve any quadratic for all its real and imaginary

roots. $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

EX: In the equation: $x^2 + 2x + 3 = 0$, find the value of the discriminant, describe the nature of the roots, then solve.

$x^2 + 2x + 3 = 0$ Determine values for a, b, and c.

$a = 1$ $b = 2$ $c = 3$ Find discriminant.

$D = 2^2 - 4 \cdot 1 \cdot 3$

$D = 4 - 12$

$D = -8$ *There are two imaginary roots.*

Solve: $x = \frac{-2 \pm \sqrt{-8}}{2}$

$x = \frac{-2 \pm 2i\sqrt{2}}{2}$

$x = -1 \pm i\sqrt{2}$

Find the value of the discriminant, describe the nature of the roots, then solve each quadratic. Use EXACT values.

60. $x^2 - 9x + 14 = 0$

61. $5x^2 - 2x + 4 = 0$

Discriminant = _____

Discriminant = _____

Type of Roots: _____

Type of Roots: _____

Roots = _____

Roots = _____

Long Division – can be used when dividing any polynomials.

Synthetic Division – can ONLY be used when dividing a polynomial by a linear (degree one) polynomial.

EX: $\frac{2x^3 + 3x^2 - 6x + 10}{x + 3}$

Long Division

$$\begin{array}{r} \frac{2x^3 + 3x^2 - 6x + 10}{x + 3} \\ 2x^2 - 3x + 3 + \frac{1}{x+3} \\ = x + 3 \overline{) 2x^3 + 3x^2 - 6x + 10} \\ \underline{(-) (2x^3 + 6x^2)} \\ -3x^2 - 6x \\ \underline{(-) (-3x^2 - 9x)} \\ 3x + 10 \\ \underline{(-) (3x + 9)} \\ 1 \end{array}$$

Synthetic Division

$$\begin{array}{r} \frac{2x^3 + 3x^2 - 6x + 10}{x + 3} \\ -3 \overline{) 2 \quad 3 \quad -6 \quad 10} \\ \quad \downarrow \quad -6 \quad 9 \quad -9 \\ \hline \quad 2 \quad -3 \quad 3 \quad 1 \\ = 2x - 3x + 3 + \frac{1}{x+3} \end{array}$$

Divide each polynomial using long division OR synthetic division.

62. $\frac{c^3 - 3c^2 + 18c - 16}{c^2 + 3c - 2}$

63. $\frac{x^4 - 2x^2 - x + 2}{x + 2}$